



GOOD BIOTECH CORP.

Website : www.good-biotech.com

E-mail : technical@good-biotech.com

38, 34<sup>th</sup> Road Taichung Industrial Park

Taichung 40768 Taiwan

Tel: +886-4-23596873

Fax:+886-4-23590410

ISO9001:2000/ISO13485:2003 & IVD GMP CERTIFIED

## SPECIFICATION SHEET

PRODUCT:	Duck IgY( $\Delta$ Fc) to <b>Digoxin</b> —affinity purified, BSA adsorbed
CATALOG NUMBER	D1805-2H
HOST ANIMAL:	Duck
DESCRIPTION:	IgY ( $\Delta$ Fc ) antibody to Digoxin is developed in duck by using Digoxin-BSA as immunogen. Affinity purified IgY ( $\Delta$ Fc ) is purified by Digoxin-BSA coupling chromatography, and adsorption is performed by BSA coupling column. The product is supplied as a solution in 10mM phosphate buffer, 0.14M sodium chloride, pH 7.4, containing 0.02% NaN <sub>3</sub> as a preservative.
IMMUNOGEN:	Digoxin-BSA
CONCENTRATION:	$\geq 1.0$ mg/ml, The concentration was determined by 280 nm by spectrophotometer, using 1.4 as the extinction coefficient
PURITY:	Determined by SDS-PAGE ( blue stain ) , showing that the IgY( $\Delta$ Fc) constitute was $\geq 90\%$ of total protein, and the total immunoglobulin IgY+IgY( $\Delta$ Fc) constitute was $\geq 90\%$ of total protein.
STORAGE:	For continuous use, store at 0~5°C. For extended storage: -20°C. Repeated freezing and thawing is not recommended.
SPECIFICITY:	N.D.
TITER:	ELISA titer: $\geq 50000x$ ELISA titer is defined as the dilution of antibody ( 1mg/ml ) sufficient to give a change in absorbance of <b>1.0</b> at 450nm. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.
APPLICATIONS:	Good to be used in immunoassay method e.g. TIA, ELISA, Immunoelectrophoresis, Western Blot assay, Immunodiffusion assay <i>et al.</i>
RELATIVE PRODUCTS:	Duck IgY( $\Delta$ Fc) to Digoxin—BSA adsorbed ( <b>D1805-2G</b> )

For *in vitro* research and the manufacturing of diagnostic only.