



GOOD BIOTECH CORP.

Website : www.good-biotech.com

E-mail : technical@good-biotech.com

38, 34th Road Taichung Industrial Park

Taichung 40768 Taiwan

Tel: +886-4-23596873

Fax:+886-4-23590410

ISO9001:2000/ISO13485:2003 & IVD GMP CERTIFIED

SPECIFICATION SHEET

PRODUCT:	Duck IgY(Δ Fc) to <i>Helicobacter pylori</i> —affinity purify
CATALOG NUMBER	D1905-2D
HOST ANIMAL	Duck
DESCRIPTION:	IgY (Δ Fc) to <i>H. pylori</i> is developed in duck by using <i>H. pylori</i> water extraction as immunogen. Affinity purified IgY (Δ Fc) is purified by <i>H. pylori</i> water extraction coupling chromatography. The product is supplied as a solution in 20mM MES (2- (N-Morpholino) ethanesulfonic acid) , 0.14M sodium chloride, pH 5.8, containing 0.02% NaN ₃ as a preservative.
IMMUNOGEN:	<i>Helicobacter pylori</i> water extraction
CONCENTRATION:	≥ 1.0 mg/ml, determined by absorbance at 280 nm and using 1.4 as the extinction coefficient
PURITY:	Determined by SDS-PAGE(blue stain), showing that the IgY(Δ Fc) constitute was $\geq 90\%$ of total protein, and the total immunoglobulin IgY+IgY(Δ Fc) constitute was $\geq 95\%$ of total protein.
STORAGE:	For continuous use, store at 0~5°C. For extended storage: -20°C. Repeating freezing and thawing is not recommended.
SPECIFICITY:	N.A.
TITER:	ELISA titer : $\geq 5000x$ ELISA titer is defined as the dilution of antibody (1mg/ml) sufficient to give a change in absorbance of 1.0 at 450nm. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.
APPLICATIONS:	Good to be used in immunoassay method e.g. ELISA, Western Blot assay, Immunoelectrophoresis, Immunodiffusion assay <i>et al.</i>
RELATIVE PRODUCTS:	Duck IgY(Δ Fc) to <i>Helicobacter pylori</i> (D1905-2B) Duck IgY and IgY(Δ Fc) to <i>Helicobacter pylori</i> urease (D1912-5B)

For *in vitro* research and manufacturing of diagnostic only.